arXiv:1303.4105v1 [quant-ph] 17 Mar 2013

Barut-Girardello and Gilmore-Perelomov coherent states for pseudoharmonic oscillator and their nonclassical properties: factorization method

M K Tavassoly[1,2,3,*] and H R Jalali[1]

[1] Atomic and Molecular Group, Faculty of Physics, Yazd University, Yazd, Iran

[2] Photonics Research Group, Engineering Research Center, Yazd University, Yazd, Iran

[3] The Laboratory of Quantum Information Processing, Yazd University, Yazd, Iran

[*] E-mail: mktavassoly@yazd.ac.ir

March 20, 2013

Keywords: Pseudoharmonic oscillator; Factorization method; Barut-Girardello coherent states; Gilmore-Perelomov coherent states; Nonclassical properties.

PACS: 42.50.Dv, 42.50.-p

Abstract

In this paper we try to introduce the ladder operators associated with the pseudoharmonic oscillator, after solving the corresponding Schrödinger equation by using the factorization method. The obtained generalized raising and lowering operators naturally lead us to the Dirac representation space of the system which is very easier to work with, in comparison to the functional Hilbert space. The $SU(1,1)$ dynamical symmetry group associated with the considered system is exactly established through investigating the fact that the deduced operators satisfy appropriate commutation relations. This result enables us to construct two important and distinct classes of Barut-Girardello and Gilmore-Perelomov coherent states associated with the system. Finally, their identities as the most important task are exactly resolved and some of their nonclassical properties are illustrated, numerically.

1 Introduction

Molecular vibrational and rotational spectroscopies are usually known as the important parts of molecular physics. The harmonic oscillator could be considered as a useful ground to explain the molecular vibrations, but this model is very ideal and simple and so has some particular restrictions. Strictly speaking, the molecular vibrations are really anharmonic, so if their Schrödinger equation can be exactly solved, they may provide more suitable models for the description of vibrating molecules. As some examples of anharmonic oscillator molecular potentials we may refer to the Morse [1], Pöschl-Teller [2, 3] and Mie-type [4] potentials which are

three typical model potentials which the molecular vibrations can be described satisfactorily by means of them. In this work, we want to study another anharmonic potential, the so-called pseudoharmonic oscillator, which is the sum of the harmonic oscillator and the inversely quadratic potential [5], i.e.,

$$V(x) = \frac{1}{2}m\omega^2 x^2 + \frac{\hbar^2}{2m}\frac{\alpha^2}{x^2}, \tag{1}$$

where m, ω and α respectively represent the mass of the particle, the frequency and the strength of the external field. Sometimes this system has been called isotonic potential [6, 7, 8, 9]. In fact, this potential was proposed by Goldman and Krivchenkov [10]. Some methods that have been used to solve the wave equation for the pseudoharmonic potential can be summarized as follows: Nikiforov-Uvarov method [11, 12], polynomial solution [13], exact quantization rule [14] and Laplace transform approach [15]. Moreover, obtaining the exact solutions of the Schrödinger equation for the pseudoharmonic oscillator using a new method (and in general enlarging the circle of solvable quantum systems) is one of the main problems in quantum physics [16].

In the present study we firstly want to solve the latter equation by the factorization method which simply arrives us to the exact solution of the eigenvalues and eigenfunctions of the considered system. Very recently this method is used for the construction of a new set of generalized coherent state and then the geometric phase of the states is investigated [17] (see also [18, 19, 20, 21] and references therein). Our motivation lies in the fact that, by using this method we will be able to find out the appropriate generalized creation and annihilation operators for the chosen system, by which we are simply lead to the Dirac space with its instruments (ket-bra space) provides us with a very simpler tool than the usual functional space. Consequently, we can construct two new, important and distinct classes of generalized coherent states associated with such physical system namely Barut-Girardello and Gilmore-Perelomov coherent states. After this exploration, due to the importance of the nonclassicality of quantum states in various theoretical and experimental fields of physics in recent decades [22, 23] we are naturally lead to the investigation of the nonclassically signs of our introduced states for completing the work. Moreover, prior to these examinations we will exactly establish the resolution of the identity of both classes of states, as the most important property of any generalized coherent state.

Notice that our present work can be considered in the line of the dynamical generalization of coherent states [24, 25, 26]. Also, it is worth mentioning that, the obtained Barut-Girardello and Gilmore-Perelomov coherent states also can be really categorized in the "algebraic" [27, 28, 29] and "group theoretical" [30, 31, 32] generalization of coherent states, respectively. From the physical point of view, we would like to mention that the strong relation of the laser light to the well-known standard bosonic coherent states corresponds to harmonic oscillator is well established from the sixth decades ego by the pioneering works of Glauber, Sudarshan and Klauder. Indeed, this was only after this discovery that, hard works along their generalizations as well as physical realization were began. This is why their various manners of generalizations are yet of hot importance [33, 34, 35, 36, 37]. Henceforth, apart from presenting a powerful technique which can be used for other known potentials, due to the direct relation of our introduced coherent states to a particular physical potential (which as we mentioned in the above lines, may be more suitable for the description of real physical phenomena) we hope that our work suitably concerns to physical realization, too.

As is mentioned before, this potential has received a lot of attention in recent years. In

addition, the used approach in this paper is conceptually near to the well-known SUSY quantum mechanics techniques which is an interesting approach for many authors [38, 39, 40, 41, 42, 43, 44]. Altogether, it may clearly be seen that our method to the solution of the eigenvalue problem and the from the final obtained results are essentially new. In this way, to the best of our knowledge, the approach to the treatment of this particular problem, the corresponding ladder operators, the associated coherent states of the types which are obtained, in addition to establishing the resolution of the identity for both classes of states and the related nonclassicality signs have not been appeared in the literature up to now.

The organization of the paper is as follows. In the next Section, we are going to review the factorization method briefly. This method is used to solve the Schrödinger equation for pseudoharmonic oscillator in one-dimensional space in Section 3. According to our obtained results, we construct the creation and annihilation operators, by which we study the dynamical group corresponding to this system in Section 4. Then, after producing the Barut-Girardello and Gilmore-Perelomov coherent states associated with the pseudoharmonic potential, the resolution of the identity corresponding to both classes of introduced states is established in Section 5. We demonstrate some of the nonclassical properties of the states like squeezing and sub-Poissonian statistics, numerically in Section 6. Finally, summary and conclusion are presented in Section 7.

2 The factorization method: a brief review

The factorization method is a useful technique that reduces the dynamical equation of a given system into a simple one which is easier to handle. This method was first introduced by Schrödinger [45, 46, 47] and then followed by Dirac [48] to avoid the usage of cumbersome mathematical tools and it has also been successfully applied to resolve some problems for which there exists exact solutions. This method further developed by Infeld and Hull [49] and Green [50]. The characteristics and advantages of the factorization method can be summarized as follows. First, this method applies only to the discrete energy spectra. Secondly, we can achieve immediately the desired eigenvalues and the normalized eigenfunctions from the given Hamiltonian without any need to use the traditional methods in quantum mechanics. Thirdly, it is possible to avoid deriving the normalization constant, which is sometimes difficult to obtain. The fourth property comes from the fact that we may discover the hidden symmetry of the quantum system through the establishment of an appropriate Lie algebra, which usually can be realized by means of the ladder operators. This method is a very powerful tool for solving second-order differential equations and has attracted much attention of many authors. In this method, which is closely related to SUSY quantum mechanics, the second-order differential operator H_0 (Hamiltonian) is written as the product of two first-order differential operators a_0 and $a_0^\dagger$ plus a constant E_0,

$$H_0 = a_0^\dagger a_0 + E_0. \tag{2}$$

Then, rewriting the SUSY Hamiltonians H_- and H_+ respectively as H_0 and H_1, and changing suitably the subscripts we have $H_1 = a_0 a_0^\dagger + E_0$. Also, by factorizing H_1 one has $H_1 = a_1^\dagger a_1 + E_1$, so firstly we have

$$H_1 = a_0 a_0^\dagger + E_0 = a_1^\dagger a_1 + E_1, \tag{3}$$

with the SUSY partner of H_1 as $H_2 = a_1 a_1^\dagger + E_1$. Continuing the above procedure leads to the generation of a hierarchy of Hamiltonians [51, 52] given by

$$H_{n+1} = a_{n+1}^\dagger a_{n+1} + E_{n+1} = a_n a_n^\dagger + E_n, \quad n = 0, 1, 2, 3, \ldots . \tag{4}$$

The complete set of eigenfunctions can be generated by the successive action of these operators on the key eigenfunctions, which are the exact solutions of the first-order differential equation. Then, the recurrence relation may be used to achieve all eigenvalues.
In fact, with the following theorem we can find eigenfunctions and eigenvalues [53].
Theorem: Assume that equations (2) to (4) with real constants E_n are true. Also suppose that there exists a null eigenfunction $|\xi_n\rangle$ with zero eigenvalue for each a_n, i.e.,

$$a_n|\xi_n\rangle = 0. \tag{5}$$

Then,

a) E_n is the nth eigenvalue of the Hamiltonian.

b) The corresponding eigenfunction of the Hamiltonian H is,

$$|\psi_n\rangle = c_n a_0^\dagger a_1^\dagger \ldots a_{n-1}^\dagger |\xi_n\rangle, \tag{6}$$

where c_n is a normalization constant may be determined as:

$$|c_n| = [(E_n - E_{n-1})(E_n - E_{n-2})\ldots(E_n - E_0)]^{-1/2}. \tag{7}$$

Three important worthwhile points lie in the theorem:

i) This method applies only to the discrete energy spectra.

ii) In the selection of a_n, one should choose it in such a way that E_n gets the maximum value.

iii) The theorem doesn't give us any information about the form of a_n. Moreover, evidently the form of a_n depends explicitly on the particular potential $V(x)$. Anyway, since $a_0^\dagger a_0$ is equal to H except for a constant, thus a_0 must contain a linear momentum term to be consistent with the kinetic energy part of the Hamiltonian. Accordingly, a_n is assumed to have the form

$$a_n = \frac{1}{\sqrt{2m}}(p + i f_n(x)), \tag{8}$$

where p is the momentum operator and $f_n(x)$ is a real (and differentiable) unknown function of x. This function would be such that equation (4) is satisfied.

3 Solving the Schrödinger equation of pseudoharmonic oscillator: factorization method

Now, we are going to search for the eigenenergies and eigenfunctions of a particular system, i.e. pseudoharmonic potential in one-dimensional space, by the factorization method. Let us start with the Schrödinger equation for this potential, i.e.,

$$\left(\frac{p^2}{2m} + V(x)\right)\psi(x) = E\psi(x), \tag{9}$$

where the operator p is defined as $\frac{\hbar}{i}\frac{d}{dx}$. According to the form of pseudoharmonic oscillator potential introduced in (1) and keeping in mind (8) we can guess that $f_n(x)$ can take the following form

$$f_n(x) = b_n x + \frac{c_n}{x}, \tag{10}$$

where b_n and c_n are constant values which should be chosen such that E_n gets the maximum value. Thus, a_n can be written as:

$$a_n = \frac{1}{\sqrt{2m}}\left[p + i\left(b_n x + \frac{c_n}{x}\right)\right]. \tag{11}$$

With the help of a_n and its conjugate $a_n^\dagger$, one can easily check that the following relations hold:

$$a_n^\dagger a_n = \frac{1}{2m}\left[p^2 + b_n^2 x^2 + b_n\left(2c_n + \hbar\right) + \frac{c_n}{x^2}\left(c_n - \hbar\right)\right], \tag{12}$$

$$a_n a_n^\dagger = \frac{1}{2m}\left[p^2 + b_n^2 x^2 + b_n\left(2c_n - \hbar\right) + \frac{c_n}{x^2}\left(c_n + \hbar\right)\right]. \tag{13}$$

With substituting $n = 0$ in (12) and using (4) it is required that

$$\begin{aligned}\frac{1}{2m}\left[p^2 + b_0^2 x^2 + b_0\left(2c_0 + \hbar\right) + \frac{c_0}{x^2}\left(c_0 - \hbar\right)\right] + E_0 \\ = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2 + \frac{\hbar^2}{2m}\frac{\alpha}{x^2}.\end{aligned} \tag{14}$$

The above equation will be satisfied, if

$$b_0^2 = m^2\omega^2 \quad \longrightarrow \quad b_0 = -m\omega, \tag{15}$$

$$c_0(c_0 - \hbar) = s(s+1)\hbar^2 \quad \longrightarrow \quad c_0 = (s+1)\hbar, \tag{16}$$

$$b_0(2c_0 + \hbar) + 2mE_0 = 0, \tag{17}$$

where the allowed values of b_0 in (15) can satisfy the condition (ii) of the stated theorem for which E_0 gets the maximum value. Also, note that the change of variable $s = -\frac{1}{2} + \sqrt{\alpha + \frac{1}{4}}$ in (16) has been used. With substituting (15) and (16) in (17), the maximum of the ground state energy will be achieved as:

$$E_0 = \hbar\omega\left(s + \frac{3}{2}\right). \tag{18}$$

For attaining the other eigenvalues we can use the recurrence relation (4), from which one has:

$$\begin{aligned}\frac{1}{2m}\left[p^2 + b_{n+1}^2 x^2 + b_{n+1}\left(2c_{n+1} + \hbar\right) + \frac{c_{n+1}}{x^2}\left(c_{n+1} - \hbar\right)\right] + E_{n+1} \\ = \frac{1}{2m}\left[p^2 + b_n^2 x^2 + b_n\left(2c_n - \hbar\right) + \frac{c_n}{x^2}\left(c_n + \hbar\right)\right] + E_n.\end{aligned} \tag{19}$$

The relation (19) will be satisfied, if

$$b_{n+1}^2 = b_n^2 \quad \longrightarrow \quad b_n = b_0 = -m\omega, \tag{20}$$

$$c_{n+1}(c_{n+1}-\hbar)=c_n(c_n+\hbar) \quad \longrightarrow \quad c_n=(s+n+1)\hbar, \tag{21}$$

$$b_{n+1}(2c_{n+1}+\hbar)+2mE_{n+1}=b_n(2c_n-\hbar)+2mE_n. \tag{22}$$

Finally, from (20), (21) and (22) we can derive the required eigenvalues as

$$E_n=\hbar\omega\left(2n+s+\frac{3}{2}\right), \quad n=0,1,2,\dots . \tag{23}$$

With substituting b_n and c_n, respectively from (20) and (21) in (11), $a_n(x)$ will be explicitly obtained

$$a_n(x)=\frac{1}{\sqrt{2m}}\left[p+i\left(-m\omega x+\frac{(s+n+1)\hbar}{x}\right)\right]. \tag{24}$$

The equation (23) gives all eigenvalues, if there exists an appropriate answer for any arbitrary n in (5). Thus, we need to solve the following equation

$$a_n(x)\xi_n(x)=0 \tag{25}$$

or equivalently

$$\frac{d\xi_n(x)}{dx}=\left(\frac{s+n+1}{x}-\frac{m\omega}{\hbar}x\right)\xi_n(x). \tag{26}$$

The solution of the above differential equation can be simply obtained as follows

$$\xi_n(x)=x^{s+n+1}e^{-\frac{m\omega}{2\hbar}x^2}. \tag{27}$$

By using (6) and after doing some tedious calculations, the normalized eigenfunctions of pseudoharmonic oscillator can be obtained as ($\hbar=m=\omega=1$):

$$\psi_n(x)=N_n x^{s+1}e^{-\frac{x^2}{2}}L_n^{s+\frac{1}{2}}(x^2), \tag{28}$$

where $L_n^{s+\frac{1}{2}}(x^2)$ is the associated Laguerre polynomials and the normalization constant in (28) is determined as

$$N_n=\sqrt{\frac{2\Gamma(n+1)}{\Gamma(n+s+\frac{3}{2})}}. \tag{29}$$

4 Introducing ladder operators associated with pseudoharmonic oscillator

It is remarkable that the operators a_n in (24) and its conjugate do not play the role of raising and lowering operators for the eigenstates of the pseudoharmonic potential. But, keeping in mind these operators, we want to construct a rather new pair of operators which yield our mentioned requirement. For this purpose, we introduce a new operator as follows

$$A_n\equiv-\frac{d}{dx}-x+\frac{s+n+1}{x}, \tag{30}$$

where its conjugate read as

$$A_n^\dagger \equiv \frac{d}{dx} - x + \frac{s+n+1}{x}. \tag{31}$$

By applying the operator A_n on $\psi_n(x)$ and using the following recurrence relations for the associated Laguerre polynomials [54]

$$x\frac{d}{dx}L_n^\alpha(x) = \begin{cases} nL_n^\alpha(x) - (n+\alpha)L_{n-1}^\alpha(x), \\ (n+1)L_{n+1}^\alpha(x) - (n+\alpha+1-x)L_n^\alpha(x), \end{cases} \tag{32}$$

we have

$$A_n\psi_n(x) = -\frac{n}{x}\psi_n(x) + \frac{2}{x}\sqrt{n(n+s+\frac{1}{2})}\,\psi_{n-1}(x). \tag{33}$$

Similarly, by acting $A_n^\dagger$ on $\psi_n(x)$ one has:

$$A_n^\dagger\psi_n(x) = -\frac{n+1}{x}\psi_n(x) + \frac{2}{x}\sqrt{(n+1)(n+s+\frac{3}{2})}\,\psi_{n+1}(x). \tag{34}$$

Clearly, the above operators $A_n, A_n^\dagger$ do not yet possess the role of ladder operators. Altogether, it is possible to find out the ladder operators which can regularly raise and lower any eigenfunctions of our considered system, i.e.,

$$M_\pm = \pm\frac{x}{2}\frac{d}{dx} + N + \frac{s}{2} + \frac{1}{\frac{3}{2}\mp\frac{1}{2}} - \frac{x^2}{2}, \tag{35}$$

where we have introduced the number operator N which satisfies the eigenvalue equation

$$N|n\rangle = n|n\rangle, \qquad \psi_n(x) = \langle x|n\rangle \tag{36}$$

with $\{|n\rangle\}_{n=0}^\infty$ as the eigenkets of the pseudoharmonic oscillator system. Notice that $N \neq M_+M_-$. It is easy to check that the ladder operators $M_\pm$ have the following properties

$$M_\pm|n\rangle = m_\pm|n\pm1\rangle, \qquad n = 0, 1, 2, ... \tag{37}$$

where

$$m_\pm = \sqrt{(n+\frac{1}{2}\pm\frac{1}{2})(n+s+\frac{2\pm1}{2})}. \tag{38}$$

We now study the dynamical group associated with the operators M_+ and M_-. Based on the obtained results in (37) and (38), we can calculate the following commutator:

$$[M_-, M_+]|n\rangle = 2m_0|n\rangle, \tag{39}$$

where we have defined

$$m_0 = n + \frac{s}{2} + \frac{3}{4}. \tag{40}$$

Thus, we can define the operator

$$M_0 = N + \frac{s}{2} + \frac{3}{4}. \tag{41}$$

Adding our above results, the operators $M_\pm$ and M_0 justify the following algebraic relations

$$[M_-, M_+] = 2M_0, \qquad [M_0, M_\pm] = \pm M_\pm. \tag{42}$$

As is well known, according to (42), the operators $M_\pm$ and M_0 correspond to an $su(1,1)$ Lie algebra. In addition, the system Hamiltonian can be simply expressed as:

$$H|n\rangle = (2n + s + \frac{3}{2})|n\rangle = [M_-, M_+]|n\rangle. \tag{43}$$

The form of the Hamiltonian in (43) satisfy the eigenvalue equation $H|n\rangle = E_n|n\rangle$ with E_n determined in (23). It may be interesting to notice that while for the harmonic oscillator one has $H_{HO}|n\rangle_{HO} = (n+1/2)|n\rangle_{HO} = \{a, a^\dagger\}|n\rangle_{HO}$, where a, $a^\dagger$ are the standard bosonic annihilation and creation operators, the anti-commutation relation for harmonic oscillator gives its place to commutation relation for pseudoharmonic oscillator.

5 Coherent states for pseudoharmonic oscillator: Barut-Girardello and Gilmore-Perelomov

We are now able to produce two distinct classes of coherent states for the considered system. Usually, the Barut-Girardello type of coherent state is defined as the right eigenstate of the annihilation operator, i.e.,

$$M_-|z\rangle_{BG} = z|z\rangle_{BG}. \tag{44}$$

To calculate the explicit form of $|z\rangle_{BG}$ with the traditional method, we write

$$|z\rangle_{BG} = \sum_{n=0}^{\infty} c_n|n\rangle. \tag{45}$$

Setting it in (44) and making use of (37) finally arrives us at the explicit form of coherent state

$$|z\rangle_{BG} = N_{BG}(|z|^2) \sum_{n=0}^{\infty} \left[\frac{\Gamma(s+\frac{3}{2})}{\Gamma(n+1)\Gamma(n+s+\frac{3}{2})}\right]^{1/2} z^n|n\rangle, \tag{46}$$

where the factor $N_{BG}(|z|^2)$ can be calculated by the normalization condition: $\langle z|z\rangle = 1$, i.e.

$$N_{BG}(|z|^2) = \left({}_0F_1(s+\frac{3}{2}, |z|^2)\right)^{-\frac{1}{2}}. \tag{47}$$

where ${}_0F_1(a,x)$ denotes the hypergeometric function.

On the other hand, referring the $su(1,1)$ symmetry algebra of the generators as in (42), we are able to construct the Gilmore-Perelomov coherent state associated with the pseudoharmonic potential as follows [55]:

$$\begin{aligned} |z\rangle_{GP} &= D(z)|0\rangle = \exp(\xi M_+ - \xi^* M_-)|0\rangle \\ &= \exp(zM_+)(1-|z|^2)^{M_0} \exp(-z^* M_-)|0\rangle, \end{aligned} \tag{48}$$

with $z = \frac{\xi}{|\xi|}\tanh(\xi)$. Straightforward calculations leads one to the corresponding normalized Gilmore-Perelomov coherent state, i.e.,

$$|z\rangle_{GP} = N_{GP}(|z|^2)\sum_{n=0}^{\infty}\left[\frac{\Gamma(n+s+\frac{3}{2})}{\Gamma(n+1)\Gamma(s+\frac{3}{2})}\right]^{1/2} z^n|n\rangle, \tag{49}$$

with the normalization factor

$$N_{GP}(|z^2|) = \left(1-|z|^2\right)^{\frac{s}{2}+\frac{3}{4}}. \tag{50}$$

Now, after finding out the two distinct classes of coherent states, we are ready to investigate some of the nonclassical properties which are of the most importance in the content of coherent states in quantum optics studies. But, before paying attention to this subject, we should investigate the resolution of the identity for the above introduced states. If we became successful, the introduced states can be truly named as coherent states.

Resolution of the identity: Any coherent state $|z\rangle$ should satisfy the resolution of the identity which is defined as follows

$$\int d^2z w(|z|^2)|z\rangle\langle z| = \hat{I} = \sum_{n=0}^{\infty}|n\rangle\langle n| \tag{51}$$

where $w(|z|^2)$ is a positive weight function that we search for obtaining it. For the first class of states which is introduced in this section, substituting (46) in (51) and using the change of variables $z = re^{i\theta}$ and $r^2 = x$ lead us to the following integral relation

$$\int_0^{\infty}\tilde{w}_{BG}(x)x^n dx = \frac{\Gamma(n+1)\Gamma(n+s+\frac{3}{2})}{\Gamma(s+\frac{3}{2})}, \tag{52}$$

where we have used the abbreviation $\tilde{w}_{BG}(x) = \pi w_{BG}(x)N^2_{BG}(x)$. This is indeed an inverse moment problem which can be solved by the well-known methods such as Mellin transform technique [56]. However, using the definition of Meijer's G-function, it follows that [57]

$$\begin{aligned}\int_0^{\infty} dx\; x^{k-1}\;\; G^{m,n}_{p,q}\;\left(\beta x\;\middle|\;\begin{matrix} a_1,\ ...,\ a_n,\ a_{n+1},\ ...,\ a_p \\ b_1,\ ...,\ b_m,\ b_{m+1},\ ...,\ b_q\end{matrix}\right) \\ = \quad \frac{1}{\beta^k}\frac{\prod_{j=1}^{m}\Gamma\left(b_j+k\right)\prod_{j=1}^{n}\Gamma\left(1-a_j-k\right)}{\prod_{j=m+1}^{q}\Gamma\left(1-b_j-k\right)\prod_{j=n+1}^{p}\Gamma\left(a_j+k\right)}.\end{aligned} \tag{53}$$

Comparing equations (52) and (53), it can be easily checked that the required weight function which can satisfy the integral equation (51) reads as:

$$w_{BG}(x) = \frac{{}_0F_1(s+\frac{3}{2},x^2)}{\pi\Gamma(s+\frac{3}{2})}G^{4,0}_{2,4}\left(x\;\middle|\;\begin{matrix}(0,s+1/2)\\(0,0,s+1/2,s+1/2)\end{matrix}\right) \tag{54}$$

where G is the well-known Meijer function and ${}_0F_1(a;x)$ is the generalized hypergeometric function.

About the Gilmore-Perelomov states, by substituting (49) in (51) we will have:

$$\int_0^{1}\tilde{w}_{GP}(x)x^n dx = \frac{\Gamma(n+1)\Gamma(s+\frac{3}{2})}{\Gamma(n+s+\frac{3}{2})}, \tag{55}$$

where we have used $\tilde{w}_{GP}(x) = \pi w_{GP}(x) N_{GP}^2(x)$. Similarly, in this case it is easy to check that the required weight function has the following form

$$w_{GP}(x) = \frac{\Gamma(s+\frac{3}{2})}{\pi\Gamma(s+\frac{1}{2})}(1-x)^{-2}{}_2F_1\left(0,0;s+\frac{1}{2};1-\frac{1}{x}\right), \tag{56}$$

where ${}_2F_1(a_1,a_2;x)$ is the generalized hypergeometric function.

A few words seem to be necessary about the limits of the integrals in (52) and (55). Indeed, any quantum state belongs legally to a Hilbert space if its normalization constant satisfy the inequality $0 < N < \infty$. This led us to the unit disk and whole of the complex plane respectively for Barut-Girardello and Gilmore-Perelomov state.

6 Nonclassical features of the introduced states

In this section, we briefly discuss some of the criteria of nonclassicality which are usually used in the relevant literature [58], and will be helpful for investigating the nonclassicality exhibition of our introduced states in the continuation of the paper. To achieve this aim, we refer to the normal squeezing, amplitude-squared squeezing and sub-Poissonian statistics as the most popular nonclassicality signs. To mention the common feature of the above criteria we should indicate that if a state is squeezed or its quantum statistics is sub-Poissonian in a region of allowed space, the corresponding Glauber-Sudarshan P-function will not be positive definite, i.e., it has no classical analogue. But, we would like to imply that finding this function is usually a hard task to do. Altogether, each of the above effects is in fact a sufficient (not necessary) condition for a quantum state to belong to the family of nonclassical states.

- *Normal squeezing:* In order to examine the quantum fluctuations of the quadratures of the field [59] we introduce the Hermitian operators

$$X_1 = \frac{M_- + M_+}{2}, \qquad P_1 = \frac{M_- - M_+}{2i}. \tag{57}$$

The squeezing parameters can be defined as follows

$$S_\gamma = \frac{(\Delta\gamma)^2}{\sqrt{\frac{1}{4}|\langle[X_1,P_1]\rangle|^2}} - 1, \;\; \gamma = X_1, P_1, \tag{58}$$

where $(\Delta\gamma)^2 = \langle\gamma^2\rangle - \langle\gamma\rangle^2$. Using (57) in (58) we obtain the following first-order squeezing parameters:

$$\begin{aligned} S_{X_1} &= \frac{1}{2\langle M_0\rangle}(\langle M_-^2\rangle + \langle M_-M_+\rangle + \langle M_+M_-\rangle + \langle M_+^2\rangle \\ &- \langle M_-^2\rangle - \langle M_+^2\rangle - \langle M_-\rangle\langle M_+\rangle - \langle M_+\rangle\langle M_-\rangle) - 1 \end{aligned} \tag{59}$$

$$\begin{aligned} S_{P_1} &= \frac{1}{2\langle M_0\rangle}(-\langle M_-^2\rangle + \langle M_-M_+\rangle + \langle M_+M_-\rangle - \langle M_+^2\rangle \\ &+ \langle M_-^2\rangle + \langle M_+^2\rangle - \langle M_-\rangle\langle M_+\rangle - \langle M_+\rangle\langle M_-\rangle) - 1, \end{aligned} \tag{60}$$

where all expectation values can be straightforwardly obtained with respect to both classes of states. A state is squeezed in X_1 or P_1 if it satisfies the inequalities $-1<S_{X_1}<0$ or $-1<S_{P_1}<0$, respectively.
By comparing the two displayed diagrams in figure (1) it can be realized that the squeezing of Gilmore-Perelomov coherent states occurs only in P_1 direction in all real space. Our calculations about the squeezing for Barut-Girardello coherent states in X_1 and P_1, show that this parameter is approximately 10^{-15} ($\cong 0$) for both operators. Thus, in view of squeezing property, it is deduced that the Barut-Girardello states associated with the considered system behave similar to the linear harmonic oscillator.

- *Amplitude-squared squeezing:* Following the proposal of Hillery [60], to evaluate this parameter we define the Hermitian operators

$$X_2=\frac{M_-^2+M_+^2}{2},\qquad P_2=\frac{M_-^2-M_+^2}{2i}. \tag{61}$$

The squeezing conditions in X_2 or P_2 are respectively given by $-1<S_{X_2}<0$ or $-1<S_{P_2}<0$, where S_{X_2} and S_{P_2} defined as follows:

$$S_\ell=\frac{(\Delta\ell)^2}{\sqrt{\frac{1}{4}|\langle[X_2,P_2]\rangle|^2}}-1,\quad \ell=X_2,P_2 \tag{62}$$

with the definition $(\Delta\ell)^2=\langle\ell^2\rangle-\langle\ell\rangle^2$. The explicit form of the latter parameters for second-order squeezing can also be obtained in a similar manner to what is done in equations (59) and (60) for the normal squeezing. In figure 2 amplitude-squared squeezing has been plotted against real z for Gilmore-Perelomov coherent states, in X_2 and P_2 direction. According to the diagrams have been placed in figure 2, it appears that, like the first-order squeezing, for this state again amplitude-squared squeezing occurs only in P_2 direction. Also, this parameter for Barut-Girardello states in the two directions X_2 and P_2 takes approximately the value of zero. This latter observation is also the same as the situation which is likely occurred in ordinary coherent states.

- *Mandel parameter:* To establish the quantum statistical properties and as another nonclassicality sign, we calculate the Mandel parameter [61], which can be generalized as:

$$Q=\frac{\langle M_+^2M_-^2\rangle-\langle M_+M_-\rangle^2}{\langle M_+M_-\rangle}-1. \tag{63}$$

The state for which $Q=0$, $Q<0$ and $Q>0$ respectively corresponds to the Poissonian (standard coherent states), sub-Poissonian (nonclassical states) and super-Poissonian (classical states) statistics. In figures 3 Mandel parameter has been plotted versus real z for Gilmore-Perelomov coherent states. As is shown, these states behave nonclassically features in a finite region of space. Our numerical calculations of this parameter for Barut-Girardello states takes the fixed value $Q=-1$. Therefore, their quantum statistics is such that Mandel parameter gets the lowest possible negative value, like the number states of harmonic oscillator which sometimes have been called as the most nonclassical states.

Adding the above observations, we may conclude that, from the point of view of squeezing properties the pseudoharmonic oscillator Barut-Girardello coherent state which is a particular superposition of number states (of pseudoharmonic oscillator) behaves like the canonical coherent states $|z\rangle = \exp(-|z|^2/2) \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{n!}} |n\rangle_{HO}$. However, taking into account the quantum statistical properties of the field, the pseudoharmonic oscillator Barut-Girardello coherent state behaves like the number states of harmonic oscillator (note that $|n\rangle_{HO} \neq |n\rangle$, where $|n\rangle$ in the present paper is related to the eigenkets of pseudoharmonic oscillator potential). So, in this view, the latter state can also be named as the most nonclassical states, too.

7 Summary and conclusion

In summary, in this paper we obtained the eigenvalues and eigenfunctions of one-dimensional pseudoharmonic oscillator by the factorization method. The presented approach allows us to find out appropriate generalized creation and annihilation operators by which one can raise and lower arbitrary eigenkets of the considered system. Then, we have explored that the system under consideration possesses a symmetrical realization of a particular dynamical group; it is shown that the constructed ladder operators can constitute the generators of the $su(1,1)$ Lie algebra. Next, we have produced the explicit form of Barut-Girardello and Gilmore-Perelomov coherent states for the considered system. The resolution of the identity, as the most important task for establishing any state as a class of "coherent states" is established analytically and so the over-completeness relation for both states are demonstrated. Finally, we could highlighted some of the nonclassical properties, such as first-order, amplitude-squared squeezing and sub-Poissonian statistics for the introduced states, numerically. Interestingly, we have also found that, in particular, considering squeezing properties, the pseudoharmonic oscillator Barut-Girardello coherent state behaves such as the canonical coherent states, however taking into account the quantum statistical properties, the latter state behaves like the number states of harmonic oscillator. Although, we have applied the method to pseudoharmonic oscillator in this paper, the approach can be used for other systems of interest, too. As another example, we have also done the same procedure to the well-known problem: a quantum particle in an infinite square well potential, successfully, with the new and novelty results, which will be appeared in near future.

References

[1] Morse P M 1929 *Phys. Rev.* **34** 57

[2] Pöschl G and Teller E 1933 *Z. Phys.* **83** 143

[3] Rosen N and Morse P M 1932 *Phys. Rev.* **42** 210

[4] Agboola D 2009 *arXiv:0812.3780v3* [math-ph] 4000

[5] Dong S H, Lozada-Cassou M, Yu J, Jimenez-Angeles F and Rivera A L 2006 *Int. J. Quan. Chem.* **107** 366

[6] Landau L D and Lifshitz E M 1977 *Quantum Mechanics: Non-Relativistic Theory 3rd edn* (New York: Pergamon) p. 127

[7] Hall R L, Saad N and von Keviczky A 2002 *J. Math. Phys.* **43** 94

[8] HallL R L and Saad N 2000 *J. Phys. A: Math. Gen.* **33** 569

[9] Saad N, Hall R L and von Keviczky A 2003 *J. Phys. A: Math. Gen.* **36** 487

[10] Goldman I I and Krivchenkov V D 1961 *Problems in Quantum Mechanics* (Pergamon, Oxford)

[11] Cheng Y F and Dai T Q 2007 *Phys. Scr.* **75** 274

[12] Berkdemir C, Berkdemir A and Han J 2006 *Chem. Phys. Lett.* **417** 326

[13] Ikhdair S M, Sever R and Mol J 2007 *Structure (THEOCHEM)* **806** 155

[14] Dong S H, Morales D and Ravelo J G 2007 *Int. J. Mod. Phys.* E **16** 189

[15] Altug Arda and Ramazan Sever 2012 *arXiv:1202.4268v1* [math-ph]

[16] Zarezadeh M and M K Tavassoly M K 2013 "The Solution of the Schrödinger Equation for a Particular Form of Morse Potential by Using Laplace Transform" *Chin. Phys. C* in press

[17] Fernandez D J 2012 *SIGMA* **8** 041

[18] Fernandez D J, Nieto L M and Rosas-Ortiz O 1995 *J. Phys. A: Math. Gen.* **28** 2693

[19] Fernandez D J, Nieto L M and Rosas-Ortiz O 1994 *J. Phys. A: Math. Gen.* **27** 3541

[20] Fernandez D J, Hussin V and Rosas-Ortiz O 2007 *J. Phys. A: Math. Theor.* **40** 6491

[21] Sbarba D and Hussin V 2007 *J. Math. Phys.* **48** 012110

[22] Wen-Xing Y, Li-Xia J, Jia-Hua L etc. 2004 *Chin. Phys. Lett.* **21** 1745

[23] Xue-li L, Lei S, Xi-wen Z etc. 1998 *Chin. Phys.* B **7** 488

[24] Nieto M M and Simmons L M 1979 *Phys. Rev.* D **20** 1321

[25] Nieto M M and Simmons L M 1979 *Phys. Rev.* D **20** 1332

[26] Nieto M M and Simmons L M 1979 *Phys. Rev.* D **20** 1342

[27] Manko V I, Marmo G, Sudarshan E C G and Zaccaria F 1997 *Phys. Scr.* **55** 528

[28] Roknizadeh R and Tavassoly M K 2004 *J. Phys. A: Math. Gen.* **37** 8111

[29] Roknizadeh R and Tavassoly M K 2005 *J. Math. Phys.* **46** 042110

[30] Gilmore R 1972 *Ann. Phys.* NY **74** 391

[31] Perelomov A M 1972 *Commun. Math. Phys.* **26** 222

[32] Barut A O and Girardello L 1971 *Commun. Math. Phys.* **21** 41

[33] Ben-Yuan Z, Lei D, Yong-Fa D, Li Y and Gao-Xiang L 2012 *Chin. Phys. B* **21** 090302

[34] Shuai W, Xiao-Yan Z and Hong-Yi F 2012 *Chin. Phys. B* **21** 054206

[35] Wei W, Guo G-C 1998 *Chin. Phys. B* **7** 174

[36] Yue-Yuan W, Ji-Cheng W and Shu-Tian L 2010 *Chin. Phys. B* **19** 074206

[37] Ahmad M A, Jie L, Yan Q, Zhi-Min M, Ai-Qun M and Shu-Tian L 2007 *Chin. Phys. B* **16** 1351

[38] Nieto M M and Simmons L M 1978 *Jr., Phys. Rev. Lett.* **41** 207

[39] Fernandez D J 2006 *AIP Conf. Proc.* **809** 80

[40] Fox R F and Choi M H 2001 *Phys. Rev.* A **64** 042104

[41] Dello Sbarba L and Hussin V 2007 *J. Math. Phys.* **48** 012110

[42] Kuru S and Negro J 2008 *Annals of Phys.* **323** 413

[43] Cooper F, Khare A and Sukhatme U 1995 *Phys. Rep.* **251** 267

[44] Khare A 2005 *AIP Conf. Proc.* **744** 133

[45] Schrödinger E 1941 *Proc. Roy. Irish Acad. Sect.* A **46** 9

[46] Schrödinger E 1941 *Proc. Roy. Irish Acad. Sect.* A **46** 183

[47] Schrödinger E 1941 *Proc. Roy. Irish Acad. Sect.* A **47** 53

[48] Dirac P M 1958 *The principles of quantum mechanics* (Oxford University Press, Oxford)

[49] Infeld I and Hull T E 1951 *Rev. Mod. Phys.* **23** 21

[50] Green H S 1968 *Matrix method in quantum mechanics* (New York: Barnes and Noble)

[51] Sukumar C 1985 *J. Phys. A: Math. Gen.* **18** L57

[52] Sukumar C 1985 *J. Phys. A: Math. Gen.* **18** 2917.

[53] Ohanian H C 1990 *Principles of quantum mechanics* (Prentice Hall. Englewood Cliffs, NJ.)

[54] Gradshteyn I S and Ryzhik I M 1994 *Tables of Integrals, Series, and Products 5th ed.* (New York: Academic Press)

[55] Miry S R and Tavassoly M K 2012 *Phys. Scr.* **85** 035404

[56] Klauder J R, Penson K A and Sixdeniers J-M 2001 *Phys. Rev.* A **64** 013817

[57] Mathai A M and Saxena R K 1973 *Generalized Hypergeometric Functions with Applications in Statistics and Physicsal Sciences* (Lecture Notes in Mathematics vol 348 Berlin: Spinger)

[58] Guang-Can G, L Jiu-Qing L 1994 Chin. Phys. B **3** 92

[59] Walls D F 1983 *Nature* **306** 141

[60] Hillery M 1987 *Opt. Commun.* **62** 135


[61] Mandel L 1979 *Opt. Lett.* **4** 205


Figure Captions:


Fig. 1 Plot of squeezing parameters, S_{X_1} (up diagram) and S_{P_1} (down diagram) against real z with $s = 1$ for Gilmore-Perelomov coherent states.

Fig. 2 Plot of amplitude-squared squeezing parameters, S_{X_2} (up diagram) and S_{P_2} (down diagram) against real z with $s = 1$ for Gilmore-Perelomov coherent states.

Fig. 3 Plot of Mandel parameter with $s = 1$ for Gilmore-Perelomov coherent states versus real z.


4
3
2
1
0
−1
S_{X_1}, S_{P_1}
−0.5
0.0
0.5
z


4
3
2
1
0
−1
S_{X_2}, S_{P_2}
−0.5
0.0
0.5
z


5
4
3
Q
2
1
0
−1
−0.5
0.0
0.5
z